arXiv:astro-ph/0211568v1 26 Nov 2002

Properties of tangential and radial angles of muons in EAS *

J. Zabierowski[a] †, K. Daumiller[b] and P. Doll[c]

[a]Soltan Institute for Nuclear Studies, Cosmic Ray Phys. Dept.,
P.O.Box 447, PL-90-950 Lodz, Poland

[b]Institut für Experimentelle Kernphysik, Universität Karlsruhe,
P.O.Box 3640, D-76021 Karlsruhe, Germany

[c]Institut für Kernphysik, Forschungszentrum Karlsruhe,
P.O. Box 3640, D-76021 Karlsruhe, Germany

Tangential and radial angles of muons in EAS, a useful concept in investigation of the muon production height, can be used also for the investigation of the muon momenta. A parameter ζ, being a combination of tangential and radial angles, is introduced and its possible applications in investigation of muons in showers are presented.

1. Introduction

EAS experiments, combining field detector arrays - being capable in measuring the shower direction - with muon tracking detectors, can be used for determination of the mean muon production height (MPH), a parameter sensitive to the primary mass. Muon tracks in air shower are, in general, not coplanar with the shower axis, which have led to the concept of tangential and radial angles [1]. However, as it will be shown in the following, these two independent observables may provide a possibility to investigate the muon momentum components in showers. A good example of the EAS experiment, where the ideas presented in this paper can be used is KASCADE [2] with its recently introduced large Muon Tracking Detector [3]. We are using data simulated with CORSIKA [4] ver.5.644 and QGSJet hadronic interaction model for all discussions and examples given below.



Figure 1. Definitions of tangential and radial angles.

2. Tangential and radial angles

As it is shown in Fig. 1, shower direction and the location of the muon detector define two perpendicular planes: *tangential* and *radial* ones.

Tangential angle τ and radial angle ρ are angles between the shower direction and the orthogonal projections of the muon track onto the tangential and radial planes, respectively.

The measured value of τ reflects the amount of muon scattering in the atmosphere and any relevant absorber/detector material, together with a possible displacement of the muon production

*Presented at XIIth ISHVECRI, CERN 2002

†e-mail: janzab@zpk.u.lodz.pl

place from the shower axis. Its distribution (Fig. 2) is symmetrical around zero and gets narrower with the increase of the muon momentum, what one could expect from the quantity related to the multiple scattering.



Figure 2. Dependence of the tangential angle on muon momentum.



Figure 3. Radial angle distribution.

A typical distribution of the radial angle is shown in Fig. 3. Its value depends on the muon parent's (meson's) transverse momentum and the distance of the muon to the shower axis at the observation level. In real experimental conditions muons are sampled within certain range of distances from the core. The shaded distribution is shown for a typical distance 20 m - 180 m from the core. The large ρ values are removed.

3. Parameter ζ

As it is seen from Fig. 1, using simple trigonometry and substituting tangent for an angle, what is correct within 5% error up to $0.4 rad$, one can write the following expressions for τ and ρ:

$$\tau \cong \frac{PB}{PD} = \frac{p_t}{p_\parallel} \times \sin \angle POB$$
$$\rho \cong \frac{PA}{PD} = \frac{p_t}{p\parallel} \times \cos \angle POB$$

Let us define the new parameter ζ as follows:

$$\varsigma \equiv \sqrt{\tau^2 + \rho^2} = \frac{p_t}{p\parallel} \ ;$$

which is valid for $\tau \leq 0.4$ rad, and $\rho \leq 0.4$ rad.

So, measuring in an experiment angles of a shower and of muons and calculating ρ and τ, by means of ζ one gets the possibility to investigate the muon momentum space. The precision of ζ depends on the precision of angular measurements in the air shower, and, in particular, on the precision of muon tracking. It is known, that the tracking accuracy improves when the high energy muons only are considered. Therefore, one would like to have as large content of high energy muons in the analyzed data sample as possible.



Figure 4. Muon momentum distribution in showers initiated by proton primaries with the energy E_0=10^{15} eV. The shaded area shows the effect of the experimental threshold 1GeV/c and the limited distance to the shower core: 20 - 180 m.

A typical muon momentum distribution in showers initiated by proton primaries with E_0=10^{15} eV is shown in Fig. 4. In all simulated showers (solid line) $\approx$ 30% are muons with momentum larger than 10 GeV/c. In real experimental conditions, where a threshold, eg.

1 GeV/c, is present and muons are registered in a restricted range of distances to the shower core, one gets the shaded distribution in Fig. 4. In this sample, with significantly reduced statistics, there is already $\approx$ 45 % of muons with p_μ $\geq$ 10 GeV/c. However, one would like to increase high energy muon content in the data sample even more, and for this purpose the ζ parameter can be used.

Indeed, as seen from Fig. 5, ζ shows strong dependence on the muon momentum. While at low momenta the full spread of ζ values is observed, its distribution becomes more confined to the low values with the increase of the muon momentum.



Figure 5. Dependence of ζ on the muon momentum. In the lower panel a mean value of ζ is plotted.

The mean value of ζ decreases with momentum exponentially, what is shown in the lower panel of Fig. 5. So, it is possible to use in the analysis the cut on the ζ value and, in this way, to increase the high energy muon content in the analyzed sample.

In Fig. 6 the ratio of muons with momentum larger than 10 GeV/c to all muons in a sample (above the threshold of 1 GeV/c) as a function of the applied cut on ζ value is plotted for two primary proton energies. Only muons in the distance between 20 and 180 m to the shower core were considered. It is seen that choosing for the analysis muons e.g. with $\zeta < 0.02$ results in significant increase of the high energy muon content in the data sample.



Figure 6. Relative content of muons with momentum larger than 10 GeV/c in the whole data sample (muons above 1 GeV/c) as a function of applied ζ cut.

Another application of the ζ parameter is rather straightforward. Its value is nothing else but a tangent of the muon angle versus the shower axis. So, knowing the distance to the shower core a simple triangulation allows to reproduce the muon production height. It is, off course, assumed that muons stem from the shower axis, what is generally not true. But, it turns out, that such approximation works well.

In Fig. 7 the true muon production height distributions (z_0) are compared with the calculated ones (MPH), using the just mentionned procedure, for two primary species, H and Fe with energy E_0=10^{15} eV. In these calculations a cut on $\zeta < 0.02$ was applied. One sees a good agreement of both curves. The mean production height values are separated by more than 2 kilometers for proton and iron. So, one can use this procedure of calculating mean muon production height for the primary mass determination.



Figure 7. Comparison of the true (z_0) and reconstructed, using parameter ζ, (MPH) muon production height distributions for two primaries, proton and iron, of E_0=10^{15} eV. Data sample was enriched in high energy muons by selecting muons with $\zeta < 0.02$.

Parameter ζ can be used for investigation of muon rapidities in showers. Substituting momentum for the total energy, which is allowed above 1 GeV without signifcant error, one can express rapidity y and pseudorapidity η using parameter ζ as follows:

$$y = \frac{1}{2}\ln\frac{E+p_\parallel}{E-p_\parallel} \approx \frac{1}{2}\ln\frac{(\sqrt{\zeta^2+1})+1}{(\sqrt{\zeta^2+1})-1}$$

$$\eta = \ln\frac{2\times p_\parallel}{p_t} \approx -\ln\frac{\zeta}{2}$$

As an example in Fig. 8 pseudorapidiy distribution of muons in showers initiated by proton primary with the energy E_0=10^{16} eV is shown. Only muons with momentum larger than 1 GeV/c are considered. Dotted line shows all pseudorapidity values for all simulated muons. Solid line corresponds to the case, where η can be calculated using parameter ζ, i.e. for muons with absolute values of τ and ρ less than 0.4 radian. Dashed line corresponds to the case, where the distance of muons to the shower core is in the range of 20 - 180 m. The large values, being the most interesting ones, are cut away due to 20 m minimum distance, so it is important to come with measurements as close to the core as possible.



Figure 8. Muon pseudorapidity distribution. For explanations see text.

The above example shows, that by means of ζ it is possible to investigate rapidities of muons in showers. In general, measuring directional parameters of muons in showers by means of the ζ parameter one can investigate muon momentum space.

The authors acknowledge very much support for this work obtained from Polish State Committee for Scientific Research (grant No. 5 P03B 133 20) and from the German Federal Ministry of Research (05 CU1VK1/9, WTZ POL 99/005).

REFERENCES

1. K. Bernlöhr, Astropart. Phys. 5 (1996) 139.
2. H.O. Klages et al.-KASCADE collaboration, Nucl. Phys. B (Proc. Suppl.) 52B (1997) 92.
3. P. Doll et al., Nucl. Instr. and Meth. A488 (2002) 517.
4. D. Heck et al., FZKA report 6019, Forschungszentrum Karlsruhe (1998).